

On Units for $B(E2)$'s

L. Zamick¹

¹*Department of Physics and Astronomy,
Rutgers University, Piscataway, New Jersey 08854, USA*

(Dated: November 17, 2009)

Abstract

In the table of Raman et. al. the units of $B(E2)$ are given as $e^2 \text{ barns}^2$ where e is the charge of the electron. Stelson and Grodzins however give $B(E2)$ in units of length^4 (cm^4).

PACS numbers:

In several different sources the formula relating the transition probability per unit time for gamma decay from a 2_1^+ state to the 0_1^+ ground state of even-even nuclei are written two different ways

$$w_1 = \frac{4\pi}{75} \frac{c}{\hbar c} \left(\frac{E}{\hbar c}\right)^5 B_1(E2; 2_1 \rightarrow 0_1) \quad (1)$$

$$w_2 = \frac{4\pi}{75} \frac{c}{\hbar c} \left(\frac{E}{\hbar c}\right)^5 e^2 B_2(E2; 2_1 \rightarrow 0) \quad (2)$$

The equivalent of w_1 is given in the works of Blatt and Weisskopf [1] and J. D. Jackson [2].

The two formulae differ in where the squared charge of the electron i.e. e^2 is placed. In w_1 the quantity e^2 is buried in $B_1(E2)$ and the dimension of B_1 is $e^2 L^4$ where L is the unit of length that is used. In w_2 the e^2 is lumped in with the other factors and the dimension of $B_2(E2)$ is just L^4 . We will use as our energy unit

MeV, and our length unit fermi. Note that $\hbar c = 197.32 \text{ MeV fm}$ and $e^2 = 1.44 \text{ MeV fm}$, while $c = 3 \times 10^{23} \text{ fm/s}$.

We then obtain:

$$w_1 = 0.85 \times 10^9 E^5 B_1(E_2) \quad (3)$$

$$w_2 = 1.23 \times 10^9 E^5 B_2(E_2) \quad (4)$$

Now in Raman's tables [3] one obtains the numerical value of the $B(E2)$ from the meanlife by using the formula

$$\tau(1 + \alpha) = 40.82 \times 10^{13} E^{-5} / (B(E2, 0 \rightarrow 2)) \quad (5)$$

Here the units are keV, barns and picoseconds where B is written (in our notation) as B_1/e^2 . But this is then simply B_2 .

Using the fact that $\tau = 1/w$, $B(E2, 0 \rightarrow 2) = 5 B(E2, 2 \rightarrow 0)$, $1 \text{ barn} = 100 \text{ fm}^2$ and $1 \text{ keV} = 10^{-3} \text{ MeV}$ we find that clearly the Raman formula is equivalent to w_2 , i.e. that the factor of e^2 is not inside $B(E2)$. In the tables of Raman et.al. the dimensions of "B(E2)" are given as $(e \text{ barn})^2$. In other words they have

B_1/e^2 in formula (5) but quote B_1 in their tables.

The expression for w_2 is given in Heyde's book in Table 4 [5]. His units for $B(E2)$, as he clearly states in eq. 4.23 and 4.27, are $(\text{fm})^4$. On the other hand, in vol. 1 of Bohr and Mottelson [6] (see eq. 3C-18), and in Lawson's book [7] (see table 5-1 and eq. 5.12) they give the numerical values of w_2 (1.23) A very simple calculation shows that the coefficient of B in their transition rate is not 1.23 but 0.85 (times powers of 10). They obviously have taken the

e^2 out of the $B(E2)$ (and multiplied this by 0.85). They perhaps confusingly use the same notation for $B_1(E2)$ and for $B(E2)$ in units of $e^2 fm^4$.

We should point out that a compilation of $B(E2)$'s similar to that of Raman was made much earlier by Stelson and Grodzins [8]. Within powers of 10 the numerical values are the same as those of Raman. Although not explicitly stated in the Stelson-Grodzins work, one can verify that they were using the w_2 relation to connect lifetimes to $B(E2)$'s. And these authors state

clearly in their work that the units of $B(E2)$ are $(\text{Length})^4$ – they use cm^4 .

-
- [1] J. M. Blatt and V. F. Weisskopf, Theoretical Nuclear Physics, John Wiley and Sons Inc., New York (1952); see Chapter XII, eq. 3.21.
 - [2] J. D. Jackson, Classical Electrodynamics, John Wiley and Sons, New York (1999); see section 9.11, eq. 9.173.
 - [3] S.Raman, C.W. Nestor, Jr. and P. Tikkanen, At. Data Nucl. Data Tables 78, 1 (2001).
 - [4] Tables of Isotopes, 8th edition, edited by R. B. Firestone and V. B. Shirley, Lawrence Berkeley Lab, John Wiley and Sons, Inc.; Appendix I-1.
 - [5] K. Heyde, The Nuclear Shell Model, Springer-Verlag, Berlin (1990).
 - [6] A. Bohr and B.R. Mottelson, Nuclear Structure, Vol. 1, W. A. Benjamin, Inc., New York, 1969.
 - [7] R. D. Lawson, Theory of the Nuclear Shell Model, Clarendon Press, Oxford, 1980.
 - [8] P. H. Stelson and L. Grodzins, Nucl. Data 1 (1965), 21.